

Superconvergence of local discontinuous Galerkin methods with generalized alternating fluxes for 1D linear convection-diffusion equations

Xiaobin Liu¹, Dazhi Zhang¹, Xiong Meng^{2,*} & Boying Wu²

¹*School of Mathematics, Harbin Institute of Technology, Harbin 150001, Heilongjiang, China;*
²*School of Mathematics and Institute for Advanced Study in Mathematics,
 Harbin Institute of Technology, Harbin 150001, Heilongjiang, China;*
Email:liuxiaobin@hit.edu.cn, zhangdazhi@hit.edu.cn, xiongmeng@hit.edu.cn, mathwby@hit.edu.cn

Received ; accepted

Abstract This paper investigates superconvergence properties of the local discontinuous Galerkin methods with generalized alternating fluxes for one-dimensional linear convection-diffusion equations. By the technique of constructing some special correction functions, we prove the $(2k + 1)$ th order superconvergence for the cell averages, and the numerical traces in the discrete L^2 norm. In addition, superconvergence of order $k + 2$ and $k + 1$ are obtained for the error and its derivative at generalized Radau points. All theoretical findings are confirmed by numerical experiments.

Keywords local discontinuous Galerkin method, superconvergence, correction function, Radau points

MSC(2010) 65M12, 65M60

Citation: Xiaobin Liu, Dazhi Zhang, Xiong Meng, Boying Wu. Superconvergence of local discontinuous Galerkin methods with generalized alternating fluxes for 1D linear convection-diffusion equations.

1 Introduction

In this paper, we consider the local discontinuous Galerkin (LDG) methods for one-dimensional linear convection-diffusion equations

$$u_t + u_x - u_{xx} = 0, \quad (x, t) \in [0, 2\pi] \times (0, T], \quad (1.1a)$$

$$u(x, 0) = u_0(x), \quad x \in R, \quad (1.1b)$$

where u_0 is sufficiently smooth. We will consider the periodic boundary condition $u(0, t) = u(2\pi, t)$, the mixed boundary condition $u(0, t) = g_1(t), u_x(2\pi, t) = g_2(t)$ and the Dirichlet boundary condition $u(0, t) = g_3(t), u(2\pi, t) = g_4(t)$. We study the superconvergence property concerning Radau points, cell averages, supercloseness of the LDG method with generalized alternating numerical fluxes, **including the case for which the parameters involved in the numerical fluxes for the prime variable regarding the convection part and the diffusion part are independently chosen for solving (1.1).**

As an extension of discontinuous Galerkin (DG) method for solving first order hyperbolic equations, the LDG method was proposed by Cockburn and Shu [16] in the framework of solving second-order convection-diffusion equations. The idea of the LDG methods is to rewrite the original equation with

*Corresponding author

high spatial derivatives as a first order system so that the DG method can be applied. Remark that in addition to stability issue, the local solvability of auxiliary variables introduced should also be guaranteed when choosing numerical fluxes.

Being a deeper insight of DG methods, superconvergence has been investigated basically measured in the discrete L^2 norm for Radau points as well as cell averages, in the L^2 norm for the error between the numerical solution and a particular projection of the exact solution (supercloseness), and in weak negative-order norm for enhancing accuracy. For example, by virtue of the duality argument in combination with the standard optimal *a priori* error estimates, Cockburn *et al.* [15] proved that the post-processed error is of order $2k + 1$ superconvergence in the L^2 norm for linear hyperbolic systems and Ji *et al.* [21] demonstrated that the smoothness-increasing accuracy-conserving (SIAC) filter can be extended to the multidimensional linear convection-diffusion equation in order to obtain $(2k + m)$ th order superconvergence, where $m = 0, \frac{1}{2}$ or 1. Here and below, k denotes the polynomial degree of the discontinuous finite element space. Later, to efficiently compute multi-dimensional problems, the line filter, namely the one-dimensional kernel is designed via rotation in [17], and a rigorous proof of the post-processed errors is also given. For arbitrary non-uniform regular meshes, by establishing the relation of the numerical solution and auxiliary variable as well as its time derivative, superconvergence of order $k + 3/2$ is proved for linear convection-diffusion equations [14]. For supercloseness results concerning high order equations, see, for example, [20, 24]. Note that aforementioned supercloseness results are not sharp. In view of this, Yang and Shu [25] adopted the dual argument to study the sharp superconvergence of the LDG method for one-dimensional linear parabolic equations, and improved superconvergence results of order $k + 2$ were obtained in terms of supercloseness and Radau points.

Recently, motivated by the successful applications of correction functions to finite element methods and finite volume methods for elliptic equations [11], Cao *et al.* [5–8] studied superconvergence properties of DG and LDG methods for linear hyperbolic and parabolic equations. Specifically, they offered a novel proof to derive the $(2k + 1)$ th or $(2k + 1/2)$ th order superconvergence rate for the cell average and numerical fluxes, which will lead to the sharp $(k + 2)$ th order superconvergence for supercloseness as well as the function errors at downwind-biased points. Note that these superconvergent results are based on a supercloseness property of the DG solution to an interpolation function consisting of the difference between a standard Gauss–Radau projection of the exact solution and a carefully designed correction function. It is worth pointing out that a suitable correction is introduced to balance the difference between the projection error for the inner product term and for the DG operator term, and in standard optimal error estimates when a Gauss–Radau projection is used, the projection error involved in the DG operator term is exactly zero. This indicates that the standard Gauss–Radau projection is not the best choice for superconvergence analysis. The superconvergence of the direct DG (DDG) method for the one-dimensional linear convection-diffusion equation was studied in [4]. We would like to remark that all the work mentioned above are focused on purely upwind and alternating numerical fluxes.

In order to obtain flexible numerical dissipation with potential applications to nonlinear systems, the upwind-biased flux was proposed in [23], which is a linear combination of the numerical solution from both sides of interfaces. Stability and optimal error estimates were obtained by constructing and analyzing some suitable *global* projections with emphasize on analysis of some circulant matrices. Note that the design of *global* projections was similar to those in the work for the Burgers–Poisson equation [22]. Moreover, Cheng *et al.* [13] studied the LDG methods for the linear convection-diffusion equations when the generalized alternating fluxes were used, and they obtained the optimal L^2 norm error estimate in a unified setting, especially when numerical fluxes with different weights are considered. In [3], Cao *et al.* investigated the superconvergence of DG methods based on upwind-biased fluxes for one-dimensional linear hyperbolic equations. More recently, Frean and Ryan [18] proved that the use of SIAC filters was still able to extract the superconvergence information and obtain a globally smooth and superconvergent solution of order $2k + 1$ for linear hyperbolic equations based on upwind-biased fluxes. Moreover, the $\alpha\beta$ -fluxes, which were introduced as linear combinations of average and jumps of the solution as well as the auxiliary variables at cell interfaces, has been a hot research topic in recent years [1, 12, 19].

In current paper, we aim at analyzing the superconvergence properties of LDG methods using gener-

alized alternating numerical fluxes for the convection-diffusion equations. The contribution of this paper is to consider the more flexible generalized alternating fluxes. The critical step in deriving superconvergence is to construct special interpolation functions for both variables (the exact solution u and the auxiliary variable q) with the aid of some suitable correction functions, essentially following [3]. Taking into account the stability result, we use special projections to eliminate or control the troublesome jump terms involving projection errors; see, e.g. Lemma 3.2. To be more precise, we will establish the superconvergence between the LDG solution (u_h, q_h) and special interpolation functions $u_I^\ell = P_\theta u - W_u^\ell$ as well as $q_I^\ell = P_\theta q - W_q^\ell$, where W_u^ℓ and W_q^ℓ are correction functions to be specified later, with the main technicality being the construction and analysis of some suitable projections tailored to the very choice of the numerical fluxes. By a rigorous mathematical proof, we prove a superconvergence rate of $2k + 1$ for the errors of numerical traces and for the cell averages, and $k + 2$ for the DG error at generalized Radau points.

The paper is organized as follows. In section 2, we present the LDG method with generalized alternating fluxes. In section 3, we construct special functions to correct the error between the LDG solution and the standard Gauss–Radau projections of the exact solution. Section 4 is the main body of the paper, in which we show and prove some superconvergence phenomena for cell averages and generalized Radau points for periodic boundary conditions. Other boundary cases including the mixed boundary condition and Dirichlet boundary condition will be considered in section 5, and the choice of numerical initial discretization is also given. In section 6, we present some numerical experiments that confirm the sharpness of our theoretical results. We will end in section 7 with concluding remarks and some possible future work.

2 The LDG scheme

In this section, we present the LDG scheme with generalized alternating fluxes for the linear convection-diffusion equation (1.1). As usual, we divide the computational domain $\Omega = [0, 2\pi]$ into N cells

$$0 = x_{\frac{1}{2}} < x_{\frac{3}{2}} < \cdots < x_{N+\frac{1}{2}} = 2\pi.$$

For any positive integer r , we define $\mathbb{Z}_r = \{1, \dots, r\}$ and denote

$$x_j = \frac{1}{2}(x_{j-\frac{1}{2}} + x_{j+\frac{1}{2}}), \quad I_j = (x_{j-\frac{1}{2}}, x_{j+\frac{1}{2}}), \quad j \in \mathbb{Z}_N$$

as the cell centers and cells, respectively. Let $h_j = x_{j+\frac{1}{2}} - x_{j-\frac{1}{2}}$ be the length of the cell I_j for $j \in \mathbb{Z}_N$, and $h = \max_{1 \leq j \leq N} h_j$. We assume the partition Ω_h is quasi-uniform in the sense that there exists a constant C independent of h such that $Ch \leq h_j \leq h$. Define the finite element space

$$V_h^k = \{v \in L^2(\Omega) : v|_{I_j} \in P^k(I_j), \forall j \in \mathbb{Z}_N\},$$

where $P^k(I_j)$ is the space of polynomials on I_j of degree at most $k \geq 0$. We use

$$\bar{u}_{j+\frac{1}{2}} = \frac{1}{2}(u_{j+\frac{1}{2}}^+ + u_{j+\frac{1}{2}}^-), \quad [u]_{j+\frac{1}{2}} = u_{j+\frac{1}{2}}^+ - u_{j+\frac{1}{2}}^-$$

to denote the mean and jump of the function u at each element boundary point $x_{j+\frac{1}{2}}$, and the weighted average is denoted by

$$u_{j+\frac{1}{2}}^{(\theta)} = \theta u_{j+\frac{1}{2}}^- + \tilde{\theta} u_{j+\frac{1}{2}}^+, \quad \tilde{\theta} = 1 - \theta,$$

where $u_{j+\frac{1}{2}}^+$ and $u_{j+\frac{1}{2}}^-$ are the traces from the right and left cells, respectively.

Throughout this paper, we employ $W^{\ell,p}(D)$ to denote the standard Sobolev space on D equipped with the norm $\|\cdot\|_{W^{\ell,p}(D)}$ with $\ell \geq 0, p = 2, \infty$. For simplicity, we set $\|\cdot\|_{W^{\ell,p}(D)} = \|\cdot\|_{\ell,p,D}$ with $D = \Omega$ or I_j . The subscript D will be omitted when $D = \Omega$, and $W^{\ell,p}(D)$ can be written as $H^\ell(D)$ when $p = 2$.

In order to construct the LDG scheme, we first introduce an auxiliary variable $q = u_x$, then the problem (1.1) can be written into a first order system

$$u_t + (u - q)_x = 0, \quad q - u_x = 0, \quad (2.1)$$

where $(u - q, u)$ is the physical flux and u is the so-called prime variable. The LDG scheme is thus to find $u_h, q_h \in V_h^k$ such that for all test functions $v, \psi \in V_h^k$

$$(u_{ht}, v)_j - (u_h - q_h, v_x)_j + (\tilde{u}_h - \hat{q}_h)v^-|_{j+\frac{1}{2}} - (\tilde{u}_h - \hat{q}_h)v^+|_{j-\frac{1}{2}} = 0, \quad (2.2a)$$

$$(q_h, \psi)_j + (u_h, \psi_x)_j - \hat{u}_h\psi^-|_{j+\frac{1}{2}} + \hat{u}_h\psi^+|_{j-\frac{1}{2}} = 0. \quad (2.2b)$$

Here $(u, v)_j = \int_{I_j} uv dx$, and $\tilde{u}_h, \hat{q}_h, \hat{u}_h$ are numerical fluxes. We use the generalized alternating numerical fluxes related to arbitrary parameters λ and θ as in [13]. That is,

$$(\tilde{u}_h - \hat{q}_h, \hat{u}_h) = (u_h^{(\lambda)} - q_h^{(\theta)}, u_h^{(\theta)}). \quad (2.3)$$

Remark that the parameters in the numerical flux regarding the convection part and diffusion part can be chosen independently, and to ensure stability the weight λ should satisfy $\lambda \geq \frac{1}{2}$.

For simplicity, we introduce the notation pertaining to the DG operator

$$\mathcal{H}^1(u, q; v) = \sum_{j=1}^N \mathcal{H}_j^1(u, q; v), \quad \mathcal{H}^2(u; \psi) = \sum_{j=1}^N \mathcal{H}_j^2(u; \psi),$$

where

$$\mathcal{H}_j^1(u, q; v) = (q - u, v_x)_j - (\hat{q} - \tilde{u})v^-|_{j+\frac{1}{2}} + (\hat{q} - \tilde{u})v^+|_{j-\frac{1}{2}},$$

$$\mathcal{H}_j^2(u; \psi) = (u, \psi_x)_j - \hat{u}\psi^-|_{j+\frac{1}{2}} + \hat{u}\psi^+|_{j-\frac{1}{2}}.$$

Thus, by Galerkin orthogonality, the cell error equation can be written as

$$(e_{ut}, v)_j + (e_q, \psi)_j + \mathcal{H}_j^1(e_u, e_q; v) + \mathcal{H}_j^2(e_u; \psi) = 0, \quad \forall v, \psi \in V_h^k, \quad (2.4)$$

where $e_u = u - u_h, e_q = q - q_h$.

For optimal error estimates of the LDG scheme using the generalized numerical fluxes (2.3) solving convection-diffusion equations with periodic boundary conditions, a globally defined projection P_θ together with P_θ^- is usually needed. For $z \in H^1(\Omega_h) = \cup_{j \in \mathbb{Z}_N} H^1(I_j)$, the generalized Gauss–Radau projection $P_\theta z$ is defined as the element of V_h^k that satisfies

$$\int_{I_j} (P_\theta z - z)v_h dx = 0, \quad \forall v_h \in P^{k-1}(I_j), \quad (2.5a)$$

$$(P_\theta z)_{j+\frac{1}{2}}^{(\theta)} = (z^{(\theta)})_{j+\frac{1}{2}}, \quad \forall j \in \mathbb{Z}_N. \quad (2.5b)$$

It has been shown in [22, 23] that the projection $P_\theta z$ is well defined for $\theta \neq \frac{1}{2}$, and for $\theta = 1/2$ some restrictions on the mesh as well as polynomial degree are needed to guarantee existence and optimal approximation property of the projection [2]. Note that when the parameter θ is taken as 0 or 1, the projection P_θ reduces to the standard local Gauss–Radau projection P_h^+ or P_h^- as defined in [10]. Besides, the projection P_θ satisfies the following optimal approximation property [22, 23]

$$\|z - P_\theta z\|_{I_j} + h^{\frac{1}{2}}\|z - P_\theta z\|_{\infty, I_j} \leq Ch^{k+\frac{3}{2}}\|z\|_{k+1, \infty}, \quad (2.6)$$

where $C > 0$ is independent of h and z .

To obtain the superconvergence results, the following lemma is useful in describing correction functions.

Lemma 2.1. [3] Suppose A is an $N \times N$ circulant matrix with the first row $(\theta, (-1)^k(1 - \theta), 0, \dots, 0)$ and the last row $((-1)^k(1 - \theta), 0, 0, \dots, \theta)$, where $\theta > 1/2$. Then, for any vectors $X = (x_1, \dots, x_N)^T, b = (b_1, \dots, b_N)^T$ satisfying $AX = b$, there holds

$$|x_j| \lesssim \|b\|_\infty, \quad \forall j \in \mathbb{Z}_N.$$

3 Correction functions

In what follows, we will present the construction of correction functions. The cases for the weights of the prime variable u_h in (2.3) being the same or different are discussed in the following two subsections.

3.1 The case with $\lambda = \theta$ in (2.3)

When $\lambda = \theta$ in (2.3), to construct special interpolation functions (u_I^ℓ, q_I^ℓ) by modifying generalized Gauss–Radau projections with correction functions so that they are superclose to the LDG solution (u_h, q_h) , we start by denoting $L_{j,k}$ as the standard Legendre polynomial of degree k on the interval I_j , and assume that the function $v(x, t)$ has the following Legendre expansion. That is, on each $I_j, j \in \mathbb{Z}_N$,

$$v(x, t) = \sum_{m=0}^{\infty} v_{j,m}(t)L_{j,m}(x), \quad v_{j,m} = \frac{2m+1}{h_j}(v, L_{j,m})_j.$$

By the definition of P_θ in (2.5a), we can rewrite $P_\theta v$ into the following form

$$P_\theta v = \sum_{m=0}^k v_{j,m}(t)L_{j,m}(x) + \bar{v}_{j,k}(t)L_{j,k}(x),$$

where $\bar{v}_{j,k}$ can be determined by $(v - P_\theta v)_{j+1/2}^{(\theta)} = 0$ with

$$v - P_\theta v = -\bar{v}_{j,k}(t)L_{j,k}(x) + \sum_{m=k+1}^{\infty} v_{j,m}(t)L_{j,m}(x). \tag{3.1}$$

It follows from the orthogonality of Legendre polynomials and (2.6) that,

$$|\bar{v}_{j,k}| \lesssim \frac{2k+1}{h_j} |(v - P_\theta v, L_{j,k})_j| \lesssim h^{k+1} \|v\|_{k+1, \infty}.$$

Following [3], to balance projection errors for the inner product term and the DG operator term, we define an integral operator D_x^{-1} by

$$D_x^{-1}u(x) = \frac{1}{\bar{h}_j} \int_{x_{j-\frac{1}{2}}}^x u(\tau) d\tau, \quad \tau \in I_j,$$

where $\bar{h}_j = h_j/2$. Obviously, $u(x) = \bar{h}_j (D_x^{-1}u(x))_x$. Moreover, by the properties of Legendre polynomials, we have

$$D_x^{-1}L_{j,k}(x) = \frac{1}{2k+1}(L_{j,k+1} - L_{j,k-1})(x). \tag{3.2}$$

To clearly see how to cancel terms involving projection errors with the goal of obtaining superconvergence, we split the error e_u, e_q into two parts:

$$\begin{aligned} e_u &= u - u_h = u - u_I^\ell + u_I^\ell - u_h \triangleq \eta_u + \xi_u, \\ e_q &= q - q_h = q - q_I^\ell + q_I^\ell - q_h \triangleq \eta_q + \xi_q. \end{aligned}$$

Then error equation (2.4) becomes

$$\begin{aligned} &(\xi_{ut}, v)_j + (\xi_q, \psi)_j + \mathcal{H}_j^1(\xi_u, \xi_q; v) + \mathcal{H}_j^2(\xi_u; \psi) \\ &= -(\eta_{ut}, v)_j - (\eta_q, \psi)_j - \mathcal{H}_j^1(\eta_u, \eta_q; v) - \mathcal{H}_j^2(\eta_u; \psi). \end{aligned}$$

For periodic boundary conditions, by choosing $v = \xi_u, \psi = \xi_q$ and summing over all j , we have

$$\frac{1}{2} \frac{d}{dt} \|\xi_u\|^2 + \|\xi_q\|^2 + \left(\lambda - \frac{1}{2}\right) \sum_{j=1}^N [\xi_u]_{j+\frac{1}{2}}^2$$

$$= -(\eta_{ut}, \xi_u) - (\eta_q, \xi_q) - \mathcal{H}^1(\eta_u, \eta_q; \xi_u) - \mathcal{H}^2(\eta_u; \xi_q). \quad (3.3)$$

From the equation (3.3), we can see that in order to obtain the supercloseness properties between the numerical solution u_h and interpolation function u_I^ℓ , we need to obtain a sharp superconvergent bound for the right-hand term, essentially using the switch of the time derivative and spatial derivative through the integral operator D_x^{-1} in combination with integration by parts; see Lemma 3.2 below. Next, we show how to construct interpolation functions and estimate the right-hand side of (3.3).

To construct the interpolation functions (u_I^ℓ, q_I^ℓ) , we define a series of functions $w_{u,i}, w_{q,i} \in V_h^k, i \in \mathbb{Z}_k$ as follows:

$$(w_{u,i} - \bar{h}_j D_x^{-1} w_{q,i-1}, v)_j = 0, \quad (w_{u,i}^{(\theta)})_{j+\frac{1}{2}} = 0, \quad (3.4a)$$

$$(w_{q,i} - w_{u,i} - \bar{h}_j D_x^{-1} \partial_t w_{u,i-1}, v)_j = 0, \quad (w_{q,i}^{(\bar{\theta})})_{j+\frac{1}{2}} = 0, \quad (3.4b)$$

where $v \in P^{k-1}(I_j)$, and

$$w_{u,0} = u - P_\theta u, \quad w_{q,0} = q - P_{\bar{\theta}} q.$$

Lemma 3.1. *The functions $w_{u,i}, w_{q,i}, i \in \mathbb{Z}_k$ defined in (3.4) have the following properties*

$$\|\partial_t w_{u,i}\|_\infty \lesssim h^{k+i+1} \|u\|_{k+i+3, \infty}, \quad \|w_{q,i}\|_\infty \lesssim h^{k+i+1} \|u\|_{k+i+2, \infty}, \quad (3.5a)$$

$$(w_{u,i}, v)_j = 0, \quad (w_{q,i}, v)_j = 0, \quad \forall v \in P^{k-i-1}(I_j). \quad (3.5b)$$

Proof. The proof of this lemma is based on deriving the following expression of $w_{u,i}$ and $w_{q,i}$ in each element I_j , which can be obtained by induction. It reads,

$$w_{u,i}|_{I_j} = \sum_{m=k-i}^k \beta_{i,m}^j L_{j,m}(x), \quad w_{q,i}|_{I_j} = \sum_{m=k-i}^k \gamma_{i,m}^j L_{j,m}(x), \quad i \in \mathbb{Z}_k. \quad (3.6)$$

Step 1: When $i = 1$, by taking $v = L_{j,m}$ with $m \leq k-1$ in (3.4a) and using (3.2) together with the orthogonality property of Legendre polynomials, we obtain

$$(w_{u,1} - \bar{h}_j D_x^{-1} w_{q,0}, v) = (\beta_{1,k-1}^j L_{j,k-1} - \frac{\bar{q}_{j,k}}{2k+1} \bar{h}_j L_{j,k-1}, v) = 0.$$

Obviously, $\beta_{1,k-1}^j = \frac{\bar{q}_{j,k}}{2k+1} \bar{h}_j$, where $\bar{q}_{j,k}$ is the coefficient of the Legendre expansion for q ; see (3.1) with v replaced by q and P_θ replaced by $P_{\bar{\theta}}$. Using the fact that $(w_{u,1}^{(\theta)})_{j+\frac{1}{2}} = 0$ we have

$$\theta \beta_{1,k}^j + (-1)^k (1-\theta) \beta_{1,k}^{j+1} = (-1)^k (1-\theta) \beta_{1,k-1}^{j+1} - \theta \beta_{1,k-1}^j. \quad (3.7)$$

Then the linear system (3.7) can be written in the matrix-vector form

$$A \beta_{1,k} = b,$$

where $A = \text{circ}(\theta, (-1)^k(1-\theta), 0, \dots, 0)$ is an $N \times N$ circulant matrix, and

$$\beta_{1,k} = \begin{pmatrix} \beta_{1,k}^1 \\ \beta_{1,k}^2 \\ \vdots \\ \beta_{1,k}^N \end{pmatrix}, \quad b = \begin{pmatrix} -\theta \beta_{1,k-1}^1 + (-1)^k (1-\theta) \beta_{1,k-1}^2 \\ -\theta \beta_{1,k-1}^2 + (-1)^k (1-\theta) \beta_{1,k-1}^3 \\ \vdots \\ -\theta \beta_{1,k-1}^N + (-1)^k (1-\theta) \beta_{1,k-1}^1 \end{pmatrix}.$$

It is easy to compute the determinant of the A in the form

$$|A| = \theta^N (1 - p^N), \quad p = \frac{(-1)^k (\theta - 1)}{\theta},$$

and for $\theta \neq \frac{1}{2}$ the matrix A is always invertible. Therefore, the linear system (3.7) has the unique solution. Moreover, by Lemma 2.1, we have

$$|\beta_{1,k}^j| \lesssim \max_{1 \leq \ell \leq N} |b_\ell| \lesssim h^{k+2} \|u\|_{k+2,\infty}, \quad \forall j \in \mathbb{Z}_N.$$

Thus,

$$\|\partial_t w_{u,1}\|_{\infty, I_j} = \|\partial_t(\beta_{1,k-1}^j L_{j,k-1} + \beta_{1,k}^j L_{j,k})\|_{\infty, I_j} \lesssim h_j |\partial_t \bar{q}_{j,k}| \lesssim h^{k+2} \|u\|_{k+4,\infty}.$$

Similarly, when choosing $v = L_{j,m}$, $m \leq k - 1$ in (3.4b), we obtain

$$w_{q,1}|_{I_j} = \sum_{m=k-1}^k \gamma_{1,m}^j L_{j,m},$$

where

$$\gamma_{1,k-1}^j = \beta_{1,k-1}^j + \frac{\partial_t \bar{u}_{j,k}}{2k+1} \bar{h}_j,$$

and $\gamma_{1,k}^j$ is the solution of the following linear system

$$\tilde{A} \gamma_{1,k} = \tilde{b},$$

with $\tilde{A} = \text{circ}(\tilde{\theta}, (-1)^k(1 - \tilde{\theta}), 0, \dots, 0)$ being an $N \times N$ circulant matrix, and

$$\gamma_{1,k} = \begin{pmatrix} \gamma_{1,k}^1 \\ \gamma_{1,k}^2 \\ \vdots \\ \gamma_{1,k}^N \end{pmatrix}, \quad \tilde{b} = \begin{pmatrix} -\tilde{\theta} \gamma_{1,k-1}^1 + (-1)^k(1 - \tilde{\theta}) \gamma_{1,k-1}^2 \\ -\tilde{\theta} \gamma_{1,k-1}^2 + (-1)^k(1 - \tilde{\theta}) \gamma_{1,k-1}^3 \\ \vdots \\ -\tilde{\theta} \gamma_{1,k-1}^N + (-1)^k(1 - \tilde{\theta}) \gamma_{1,k-1}^1 \end{pmatrix}.$$

Consequently, the estimate of $\|w_{q,1}\|_{\infty}$ in (3.5a) follows by using Lemma 2.1 and optimal approximation property (2.6). Moreover, (3.5b) is a trivial consequence of the expression (3.6) when the orthogonality property of Legendre polynomials is taken into account.

Step 2: Suppose that (3.5a) and (3.6) are valid for all $i \leq k - 1$ and we want to prove it still holds for $i + 1$. From equation (3.4a), we can get

$$\left(w_{u,i+1} - \bar{h}_j D_x^{-1} \left(\sum_{m=k-i}^k \gamma_{i,m}^j L_{j,m} \right), v \right)_j = 0, \quad \forall v \in P^{k-1}(I_j).$$

In order to get the superconvergent bounds of $w_{u,i+1}$, we need to find out the expression of coefficient $\beta_{i+1,m}$. After a direct calculation, there hold

$$\beta_{i+1,k-i-1}^j = -\frac{\gamma_{i,k-i}^j \bar{h}_j}{2(k-i)+1}, \quad \beta_{i+1,k-i}^j = -\frac{\gamma_{i,k-i+1}^j \bar{h}_j}{2(k-i)+3},$$

$$\beta_{i+1,m}^j = \bar{h}_j \left(\frac{\gamma_{i,m-1}^j}{2m-1} - \frac{\gamma_{i,m+1}^j}{2m+3} \right), \quad m = k-i+1, \dots, k-1.$$

Moreover, by the fact that $w_{u,i+1}^{(\theta)} = 0$, we get

$$\theta(\beta_{i+1,k-i-1}^j + \dots + \beta_{i+1,k}^j) + (1-\theta)(-1)^{k-i-1} \beta_{i+1,k-i-1}^{j+1} + \dots + (1-\theta)(-1)^k \beta_{i+1,k}^{j+1} = 0.$$

Again, we can write the above linear system into the matrix-vector form

$$A \beta_{i+1,k} = c,$$

and when $\theta \neq \frac{1}{2}$, we arrive at the unique existence of the system. Consequently, it follows from Lemma 2.1 that

$$\begin{aligned} \|\partial_t w_{u,i+1}\|_{\infty, I_j} &\lesssim \sum_{m=k-i-1}^k |\partial_t \beta_{i+1,m}^j| \lesssim h \sum_{m=k-i}^k |\partial_t \gamma_{i,m}^j| \\ &\lesssim h \|\partial_t w_{q,i}\|_{\infty} \lesssim h^{k+i+2} \|\partial_t q\|_{k+i+1, \infty} \lesssim h^{k+i+2} \|u\|_{k+i+4, \infty}. \end{aligned}$$

Analogously, the other estimate of (3.5a) can be obtained, and the orthogonality property in (3.5b) is a trivial consequence of expression of $w_{u,i}$ and $w_{q,i}$ in (3.6) with i replaced by $i+1$. This finishes the proof of Lemma 3.1. \square

We are now ready to define the correction functions as follows. For any positive integer $\ell \in \mathbb{Z}_k$, we define in each element I_j

$$W_u^\ell = \sum_{i=1}^{\ell} w_{u,i}, \quad W_q^\ell = \sum_{i=1}^{\ell} w_{q,i}, \quad (3.8)$$

and the special interpolation functions are

$$u_I^\ell = P_\theta u - W_u^\ell, \quad q_I^\ell = P_\theta q - W_q^\ell. \quad (3.9)$$

Lemma 3.2. *Suppose $u \in W^{k+\ell+3, \infty}(\Omega)$, $\ell \in \mathbb{Z}_k$ is the solution of (1.1), and u_I^ℓ, q_I^ℓ are defined by (3.9), then for $\forall v, \psi \in V_h^k$, we have*

$$|((u - u_I^\ell)_t, v)_j - (W_u^\ell, v_x)_j + (W_q^\ell, v_x)_j| \lesssim h^{k+\ell+1} \|u\|_{k+\ell+3, \infty} \|v\|_{1, I_j}, \quad (3.10a)$$

$$|(q - q_I^\ell, \psi)_j + (W_u^\ell, \psi_x)_j| \lesssim h^{k+\ell+1} \|u\|_{k+\ell+2, \infty} \|\psi\|_{1, I_j}. \quad (3.10b)$$

Proof. By the orthogonality property of $w_{u,i}$ and $w_{q,i}$, $i \in \mathbb{Z}_{k-1}$, we have

$$\begin{aligned} D_x^{-1} w_{u,i}(x_{j+\frac{1}{2}}^-) &= \frac{1}{h_j} (w_{u,i}, 1)_j = 0 = D_x^{-1} w_{u,i}(x_{j-\frac{1}{2}}^+), \\ D_x^{-1} w_{q,i}(x_{j+\frac{1}{2}}^-) &= \frac{1}{h_j} (w_{q,i}, 1)_j = 0 = D_x^{-1} w_{q,i}(x_{j-\frac{1}{2}}^+). \end{aligned}$$

It follows from integration by parts that

$$\begin{aligned} (\partial_t w_{u,i}, v)_j &= -\bar{h}_j (D_x^{-1} \partial_t w_{u,i}, v_x)_j = -(w_{q,i+1} - w_{u,i+1}, v_x), \quad v \in V_h^k, \quad i \in \mathbb{Z}_{k-1}, \\ (w_{q,i}, v)_j &= -\bar{h}_j (D_x^{-1} w_{q,i}, v_x)_j = -(w_{u,i+1}, v_x), \quad v \in V_h^k, \quad i \in \mathbb{Z}_{k-1}. \end{aligned}$$

Then

$$\begin{aligned} &((u - u_I^\ell)_t, v)_j - (W_u^\ell, v_x)_j + (W_q^\ell, v_x)_j \\ &= ((u - P_\theta u)_t, v)_j + \sum_{i=1}^{\ell} [(\partial_t w_{u,i}, v)_j + (w_{q,i} - w_{u,i}, v_x)_j] \\ &= (\partial_t w_{u,\ell}, v)_j. \end{aligned}$$

Similarly, there holds

$$(q - q_I^\ell, \psi)_j + (W_u^\ell, \psi_x)_j = (w_{q,\ell}, \psi)_j.$$

By (3.5a), we can get the desired result (3.10). \square

3.2 The case with $\lambda \neq \theta$ in (2.3)

When parameters λ and θ in (2.3) pertaining to convection and diffusion terms are chosen differently, a pair of suitable interpolation functions in possession of supercloseness property should be constructed, which are based on a combination of modified projections and new correction functions. To do that, let us first recall a new modified projection [13]. That is,

$$\Pi_h(u, q) = (P_\theta u, P_\theta^* q),$$

in which $P_\theta u \in V_h^k$ has been given in (2.5a), and $P_\theta^* q \in V_h^k$ depends on both u and q satisfying

$$\begin{aligned} \int_{I_j} (P_\theta^* q) v_h dx &= \int_{I_j} q v_h dx, \quad \forall v_h \in P^{k-1}(I_j), \\ (P_\theta^* q)_{j+\frac{1}{2}}^{(\bar{\theta})} &= (q^{(\bar{\theta})})_{j+\frac{1}{2}} + (\lambda - \theta)[u - P_\theta u]_{j+\frac{1}{2}} \end{aligned}$$

for any $j = 1, \dots, N$. Moreover, this projection have the following approximation property

$$\|q - P_\theta^* q\|_{I_j} \leq Ch^{k+\frac{3}{2}} \left(\|q\|_{k+1, \infty} + |\lambda - \theta| \cdot \|u\|_{k+1, \infty} \right).$$

From the above estimate of $q - P_\theta^* q$, it is easy to see that the coefficient $\bar{q}_{j,k}$ can be controlled by the prime and auxiliary variables, namely

$$\begin{aligned} |\bar{q}_{j,k}| &\lesssim \frac{2k+1}{h_j} |(q - P_\theta^* q, L_{j,k})| \\ &\lesssim h^{k+1} \left(\|q\|_{k+1, \infty} + |\lambda - \theta| \cdot \|u\|_{k+1, \infty} \right) \\ &\lesssim h^{k+1} \|u\|_{k+2, \infty}. \end{aligned}$$

Next, the corresponding correction functions pertaining to two different weights λ and θ can be easily defined. Specifically, we define the functions $w_{u,i}, w_{q,i}, i \in \mathbb{Z}_k$ satisfying

$$(w_{u,i} - \bar{h}_j D_x^{-1} w_{q,i-1}, z)_j = 0, \quad (w_{u,i}^{(\theta)})_{j+\frac{1}{2}} = 0, \quad (3.11a)$$

$$(w_{q,i} - w_{u,i} - \bar{h}_j D_x^{-1} \partial_t w_{u,i-1}, z)_j = 0, \quad (w_{q,i}^{(\bar{\theta})})_{j+\frac{1}{2}} = (w_{u,i}^{(\lambda)})_{j+\frac{1}{2}}, \quad (3.11b)$$

where $z \in P^{k-1}(I_j)$, and $w_{u,0} = u - P_\theta u, w_{q,0} = q - P_\theta^* q$.

Let us finish this section by providing the following theorem.

Theorem 3.3. *When $\lambda \neq \theta$ in (2.3), the functions $w_{u,i}, w_{q,i}, i \in \mathbb{Z}_k$ defined in (3.11) still have the following properties*

$$\begin{aligned} \|\partial_t w_{u,i}\|_\infty &\lesssim h^{k+i+1} \|u\|_{k+i+3, \infty}, \quad \|w_{q,i}\|_\infty \lesssim h^{k+i+1} \|u\|_{k+i+2, \infty}, \\ (w_{u,i}, v)_j &= 0, \quad (w_{q,i}, v)_j = 0, \quad \forall v \in P^{k-i-1}(I_j). \end{aligned}$$

Moreover, when $u \in W^{k+\ell+3, \infty}(\Omega), \ell \in \mathbb{Z}_k$, the special interpolation functions

$$u_I^\ell = P_\theta u - W_u^\ell, \quad q_I^\ell = P_\theta^* q - W_q^\ell$$

with (3.8) satisfy

$$\begin{aligned} |((u - u_I^\ell)_t, v)_j - (W_u^\ell, v_x)_j + (W_q^\ell, v_x)_j| &\lesssim h^{k+\ell+1} \|u\|_{k+\ell+3, \infty} \|v\|_{1, I_j}, \\ |(q - q_I^\ell, \psi)_j + (W_u^\ell, \psi_x)_j| &\lesssim h^{k+\ell+1} \|u\|_{k+\ell+2, \infty} \|\psi\|_{1, I_j}. \end{aligned}$$

Proof. Since there is only slight difference between (3.4) and (3.11) in terms of different boundary collocations, Theorem 3.3 can thus be proved by an argument similar to that in subsection 3.1 with different vectors b, \tilde{b} and c , etc. The detailed proof is omitted. \square

4 Superconvergence

In this section, we will show the superconvergence properties for the LDG solution at some special points as well as cell averages, which are mainly based on the supercloseness result for the error between the LDG solution (u_h, q_h) and newly designed interpolation functions (u_I^ℓ, q_I^ℓ) .

Theorem 4.1. *Let $u \in W^{k+\ell+3, \infty}(\Omega)$, $\ell \in \mathbb{Z}_k$ is the exact solution of (1.1), and u_h, q_h are the numerical solutions of LDG scheme (2.2), Then for periodic boundary conditions*

$$\|u_I^\ell - u_h\| + \left(\int_0^t \|q_I^\ell - q_h\|^2 d\tau \right)^{\frac{1}{2}} \leq C(1+t)h^{k+\ell+1},$$

where C depends on $\|u\|_{k+\ell+3, \infty}$.

Proof. Using Lemma 3.2, we obtain

$$\begin{aligned} & |(\eta_{ut}, v)_j + (\eta_q, \psi)_j + \mathcal{H}_j^1(\eta_u, \eta_q; v) + \mathcal{H}_j^2(\eta_u; \psi)| \\ &= |((u - u_I^\ell)_t, v)_j - (W_u^\ell, v_x)_j + (W_q^\ell, v_x)_j + (q - q_I^\ell, \psi)_j + (W_u^\ell, \psi_x)_j| \\ &= |(\partial_t w_{u, \ell}, v)_j + (w_{q, \ell}, \psi)_j| \\ &\lesssim h^{k+\ell+1} \|u\|_{k+\ell+3, \infty} (\|v\|_{1, I_j} + \|\psi\|_{1, I_j}). \end{aligned}$$

Inserting the above estimate into (3.3) and summing over all j , one has

$$\frac{1}{2} \frac{d}{dt} \|\xi_u\|^2 + \|\xi_q\|^2 \lesssim h^{k+\ell+1} \|u\|_{k+\ell+3, \infty} (\|\xi_u\| + \|\xi_q\|).$$

If we choose a suitable initial condition such that

$$\|\xi_u(0)\| = 0,$$

then Theorem 4.1 follows by using Young's inequality and Gronwall inequality. \square

4.1 Superconvergence of numerical fluxes

In this subsection, we present the superconvergence results of the numerical fluxes.

Theorem 4.2. *Assume that $u \in W^{2k+3, \infty}(\Omega)$, $k \geq 1$ is the solution of (1.1), and the u_h, q_h are the numerical solutions of LDG scheme (2.2) with the initial solution $u_h(\cdot, 0) = u_I^k(\cdot, 0)$. Then for periodic boundary conditions*

$$\|e_{u, n}\| \lesssim C(1+t)h^{2k+1}, \quad \left(\int_0^t \|e_{q, n}\|^2 d\tau \right)^{\frac{1}{2}} \lesssim C(1+t)h^{2k+1},$$

where

$$\|e_{v, n}\| = \left(\frac{1}{N} \sum_{j=1}^N |(v - \hat{v}_h)(x_{j+\frac{1}{2}}, t)|^2 \right)^{\frac{1}{2}}, \quad v = u, q.$$

Proof. It follows from the inverse inequality and the supercloseness result in Theorem 4.1 that

$$\begin{aligned} \|e_{u, n}\| &= \left(\frac{1}{N} \sum_{j=1}^N |(\hat{u}_I - \hat{u}_h)(x_{j+\frac{1}{2}}, t)|^2 \right)^{\frac{1}{2}} \\ &\lesssim \left(\frac{1}{N} \sum_{j=1}^N h_j^{-1} \|u_I - u_h\|_{I_j \cup I_{j+1}}^2 \right)^{\frac{1}{2}} \\ &\lesssim \|u_I^k - u_h\| \lesssim C(1+t)h^{2k+1}. \end{aligned}$$

Using the supercloseness result in Theorem 4.1 again, superconvergence of the auxiliary variable q can be derived analogously. This finishes the proof of Theorem 4.2. \square

4.2 Superconvergence for cell averages

Theorem 4.3. Assume that the conditions of Theorem 4.1 are satisfied, then for the periodic boundary conditions, we have

$$\|e_u\|_c \lesssim (1+t)h^{2k+1}\|u\|_{2k+3,\infty}, \quad \left(\int_0^t \|e_q\|_c^2 d\tau\right)^{\frac{1}{2}} \lesssim (1+t)h^{2k+1}\|u\|_{2k+3,\infty}, \quad (4.1)$$

where

$$\|e_v\|_c = \left(\frac{1}{N} \sum_{j=1}^N \left(\frac{1}{h_j} (e_v, 1)_j\right)^2\right)^{\frac{1}{2}}, \quad v = u, q.$$

Proof. Taking $\|e_u\|_c$ as an example, by the properties of P_θ and the definition of u_I^k , we obtain

$$(e_u, 1)_j = (u_I^k - u_h, 1)_j + (W_u^k, 1)_j. \quad (4.2)$$

The superconvergent result can thus be proved by using the orthogonality property in (3.5b), Cauchy–Schwarz inequality and Theorem 4.1. \square

4.3 Superconvergence at generalized Radau points

As a natural extension of Radau points for $\theta = 1$, the roots of generalized Radau polynomials for the weight θ are introduced in [18]. To be more specific, the generalized Radau polynomials are defined as

$$R_{k+1}^* = \begin{cases} L_{k+1} - (2\theta - 1)L_k, & \text{when } k \text{ is even,} \\ (2\theta - 1)L_{k+1} - L_k, & \text{when } k \text{ is odd.} \end{cases} \quad (4.3)$$

For superconvergence analysis, instead of using the global projection $P_\theta u$, a much simpler local projection $P_h u$ is introduced [3]:

$$\begin{aligned} \int_{I_j} (P_h u - u)v &= 0, \quad \forall v \in P^{k-1}(I_j), \\ \theta P_h u(x_{j+\frac{1}{2}}^-) + (1-\theta)P_h u(x_{j-\frac{1}{2}}^+) &= \theta u(x_{j+\frac{1}{2}}^-) + (1-\theta)u(x_{j-\frac{1}{2}}^+). \end{aligned}$$

Lemma 4.4. [3] Suppose $u \in W^{k+2,\infty}(\Omega)$ and $P_h u$ is the local projection of u defined above with $\theta \neq \frac{1}{2}$, then

$$\begin{aligned} |(u - P_h u)(R_{j,m}^r)| &\lesssim h^{k+2}\|u\|_{k+2,\infty}, \\ |\partial_x(u - P_h u)(R_{j,m}^{r*})| &\lesssim h^{k+1}\|u\|_{k+2,\infty}, \\ \|P_h u - P_\theta u\|_\infty &\lesssim h^{k+2}\|u\|_{k+2,\infty}. \end{aligned}$$

Here $R_{j,m}^r, R_{j,m}^{r*}$ are the roots of rescaled Radau polynomials $R_{j,m+1}^*$ and $\partial_x R_{j,m+1}^*$.

We are now ready to show the superconvergence result at generalized Radau points.

Theorem 4.5. Let $u \in W^{k+5,\infty}(\Omega)$ and u_h be the numerical solution of (1.1), suppose $u_I^\ell, \ell \geq 2$ is the special interpolation function defined in (3.9), then for the periodic boundary conditions

$$\begin{aligned} \|e_{u,r}\| &\lesssim (1+t)h^{k+2}\|u\|_{k+5,\infty}, & \|e_{u,rx}\| &\lesssim (1+t)h^{k+1}\|u\|_{k+5,\infty}, \\ \left(\int_0^t \|e_{q,l}\|^2 d\tau\right)^{\frac{1}{2}} &\lesssim (1+t)h^{k+2}\|u\|_{k+5,\infty}, & \left(\int_0^t \|e_{q,lx}\|^2 d\tau\right)^{\frac{1}{2}} &\lesssim (1+t)h^{k+1}\|u\|_{k+5,\infty}, \end{aligned}$$

where

$$\begin{aligned} \|e_{u,r}\| &= \max_{j \in \mathbb{Z}_N} |(u - u_h)(R_{j,m}^r)|, & \|e_{u,rx}\| &= \max_{j \in \mathbb{Z}_N} |(u - u_h)_x(R_{j,m}^{r*})|, \\ \|e_{q,l}\| &= \max_{j \in \mathbb{Z}_N} |(q - q_h)(R_{j,m}^l)|, & \|e_{q,lx}\| &= \max_{j \in \mathbb{Z}_N} |(q - q_h)_x(R_{j,m}^{l*})|. \end{aligned}$$

Here, $R_{j,m}^l$ and $R_{j,m}^{l*}$ are the roots of rescaled Radau polynomials $R_{j,m+1}^*$ and $\partial_x R_{j,m+1}^*$ in (4.3) with θ replaced by $\tilde{\theta}$.

Proof. By choosing $\ell = 2$ in Theorem 4.1, we obtain

$$\|u_h - u_I^2\| \lesssim (1+t)h^{k+3}\|u\|_{k+5,\infty}.$$

From the inverse inequality, we can get

$$\|u_h - u_I^2\|_\infty \lesssim h^{-\frac{1}{2}}\|u_h - u_I^2\| \lesssim (1+t)h^{k+\frac{5}{2}}\|u\|_{k+5,\infty}.$$

By the the triangle inequality,

$$\begin{aligned} |(u - u_h)(R_{j,m}^r)| &\lesssim \|u_h - u_I^2\|_\infty + |(u - P_h u)(R_{j,m}^r)| + \|P_h u - P_\theta u\|_\infty + \|W_u^2\|_\infty \\ &\lesssim h^{k+2}\|u\|_{k+2,\infty}. \end{aligned}$$

The superconvergence results for the derivative of errors and the auxiliary variable q can be obtained by the same arguments. This completes the proof of Theorem 4.5. \square

Remark 4.6. The analysis of superconvergence is mainly based on the supercloseness between the LDG solution (u_h, q_h) and the interpolation function (u_I^ℓ, q_I^ℓ) by asking for (W_u^ℓ, W_q^ℓ) satisfying

$$(W_u^\ell)_{j+\frac{1}{2}}^{(\theta)} = 0, \quad (W_q^\ell)_{j+\frac{1}{2}}^{(\bar{\theta})} = 0, \quad j \in \mathbb{Z}_N.$$

Therefore, when $\lambda \neq \theta$, the superconvergence results for auxiliary variable q are no longer valid, since (3.11b) is needed indicating that $(W_q^\ell)_{j+\frac{1}{2}}^{(\bar{\theta})} \neq 0$. In addition, when $\lambda = \theta$, superconvergence of q can be proved in the $L^2([0, T]; L^2[0, 2\pi])$ norm while superconvergence can be observed numerically in the $L^\infty([0, T]; L^2[0, 2\pi])$ norm.

5 Other boundary conditions

5.1 Mixed boundary conditions

For the mixed boundary conditions

$$u(0, t) = g_1(t), \quad u_x(2\pi, t) = g_2(t), \quad (5.1)$$

the numerical fluxes are chosen as

$$(\tilde{u}_h - \hat{q}_h, \hat{u}_h)_{j+\frac{1}{2}} = \begin{cases} (g_1 - q_h^+, g_1), & j = 0, \\ (u_h^\theta - q_h^{\bar{\theta}}, u_h^\theta), & j = 1, \dots, N-1, \\ (u_h^- - g_2, u_h^-), & j = N. \end{cases} \quad (5.2)$$

The corresponding global projections P_θ and $P_{\bar{\theta}}$ are modified to be in the following piecewise global version, i.e.,

$$\begin{cases} (\tilde{P}_\theta u, v)_j = (u, v)_j, \quad \forall v \in P^{k-1}(I_j), \\ (\tilde{P}_\theta u)_{j+\frac{1}{2}}^{(\theta)} = u_{j+\frac{1}{2}}^{(\theta)}, \quad j = 1, \dots, N-1, \\ (\tilde{P}_\theta u)_{N+\frac{1}{2}}^- = u_{N+\frac{1}{2}}^-, \quad j = N, \end{cases}$$

and

$$\begin{cases} (\tilde{P}_{\bar{\theta}} q, \eta)_j = (q, \eta)_j, \quad \forall \eta \in P^{k-1}(I_j), \\ (\tilde{P}_{\bar{\theta}} q)_{j-\frac{1}{2}}^{(\bar{\theta})} = q_{j-\frac{1}{2}}^{(\bar{\theta})}, \quad j = 2, \dots, N, \\ (\tilde{P}_{\bar{\theta}} q)_{\frac{1}{2}}^+ = q_{\frac{1}{2}}^+, \quad j = 1. \end{cases} \quad (5.3)$$

Obviously, the projection \tilde{P}_θ can be decoupled starting from the cell I_N and $\tilde{P}_{\bar{\theta}}$ can be computed from the cell I_1 . Moreover, we have the following optimal approximation properties.

Lemma 5.1. [23] Assume $z \in W^{k+1,\infty}(I_j)$ with $\theta \neq \frac{1}{2}$, the projection $P = \tilde{P}_\theta$ or $\tilde{P}_{\tilde{\theta}}$ defined above satisfies the following approximation property

$$\|z - Pz\|_{I_j} + h^{\frac{1}{2}}\|z - Pz\|_{\infty, I_j} \leq Ch^{k+\frac{3}{2}}\|z\|_{k+1,\infty},$$

where C is independent of h and z .

Replacing P_θ ($P_{\tilde{\theta}}$) by \tilde{P}_θ ($\tilde{P}_{\tilde{\theta}}$), we are able to construct the following correction functions in possession of supercloseness properties. That is, for $z \in P^{k-1}(I_j)$

$$\begin{aligned} (w_{u,i} - \bar{h}_j D_x^{-1} w_{q,i-1}, z)_j &= 0, & (w_{u,i}^{(\theta)})_{j+\frac{1}{2}} &= 0, \quad \forall j \in \mathbb{Z}_{N-1}, \\ (w_{q,i} - w_{u,i} - \bar{h}_j D_x^{-1} \partial_t w_{u,i-1}, z)_j &= 0, & (w_{q,i}^{(\tilde{\theta})})_{j+\frac{1}{2}} &= 0, \quad \forall j \in \mathbb{Z}_{N-1}, \\ & & (w_{u,i}^-)_{N+\frac{1}{2}} &= 0, \quad (w_{q,i}^+)_{\frac{1}{2}} = 0. \end{aligned}$$

The superconvergent results can thus be obtained if we follow the same argument as that in section 3 and section 4.

5.2 Dirichlet boundary conditions

For Dirichlet boundary conditions

$$u(0, t) = g_3(t), \quad u(2\pi, t) = g_4(t), \tag{5.4}$$

following [9], we choose the numerical fluxes as

$$(\tilde{u}_h - \hat{q}_h, \hat{u}_h)_{j+\frac{1}{2}} = \begin{cases} (g_3 - q_h^+, g_3), & j = 0, \\ (u_h^\theta - q_h^\theta, u_h^\theta), & j = 1, \dots, N-1, \\ (u_h^- - q_h^-, g_4), & j = N. \end{cases}$$

Similarly, we still need to make slight changes to the projection. For projection $\tilde{P}_{\tilde{\theta}}$, we still adopt the definition in (5.3), while the projection \tilde{P}_θ is modified as follows:

$$\begin{cases} (\tilde{P}_\theta u, v)_j = (u, v)_j, \quad \forall v \in P^{k-1}(I_j), \\ (\tilde{P}_\theta u)_{j+\frac{1}{2}}^{(\theta)} = u_{j+\frac{1}{2}}^{(\theta)}, \quad j \in \mathbb{Z}_{N-1}, \\ (\tilde{P}_\theta u)_{N+\frac{1}{2}}^- = u_{N+\frac{1}{2}}^- + (\tilde{P}_\theta q - q)_{N+\frac{1}{2}}^-. \end{cases}$$

From the last equation we can see that, compared with the mixed boundary condition, the left limit of projection at point $x_{N+\frac{1}{2}}$ consists of two parts. One is the left limit of exact solution u at point $x_{N+\frac{1}{2}}$, the other is the left limit of projection error of the auxiliary variable q at point $x_{N+\frac{1}{2}}$. Since we do not have any information about the auxiliary variable q at the boundary, we need to use the prime variable u to eliminate the boundary term introduced by $\tilde{P}_\theta q - q$ at point $x_{N+\frac{1}{2}}$.

The superconvergent results can be obtained if we define the following correction functions: for $z \in P^{k-1}(I_j)$,

$$\begin{aligned} (w_{u,i} - \bar{h}_j D_x^{-1} w_{q,i-1}, z)_j &= 0, & (w_{u,i}^{(\theta)})_{j+\frac{1}{2}} &= 0, \quad \forall j \in \mathbb{Z}_{N-1}, \\ (w_{q,i} - w_{u,i} - \bar{h}_j D_x^{-1} \partial_t w_{u,i-1}, z)_j &= 0, & (w_{q,i}^{(\tilde{\theta})})_{j+\frac{1}{2}} &= 0, \quad \forall j \in \mathbb{Z}_{N-1}, \\ & & (w_{q,i}^+)_{\frac{1}{2}} &= 0, \quad (w_{u,i}^-)_{N+\frac{1}{2}} = (w_{q,i}^-)_{N+\frac{1}{2}}. \end{aligned}$$

5.3 Initial discretization

In this section, we consider how to discretize the initial datum. Initial value discretization is very important for the study of superconvergence, which can be obtained using the same technique as that in [3]. Specifically, for periodic boundary conditions,

1. according to the definition of projection $P_\theta, P_{\bar{\theta}}$, calculate the $w_{u,0}, w_{q,0}$;
2. calculate $w_{u,i}, w_{q,i}$ by the equations (3.4);
3. calculate $W_u^\ell = \sum_{i=1}^{\ell} w_{u,i}, u_I^\ell = P_\theta u - W_u^\ell$;
4. let $u_h(\cdot, 0) = u_I^\ell(\cdot, 0)$.

6 Numerical results

In this section, we provide numerical examples to illustrate our theoretical findings. For time discretization, we use TVDRK3 method and take $\Delta t = CFL * h^2$.

Example 6.1. We consider the following problem

$$\begin{aligned} u_t + u_x - u_{xx} &= 0, & (x, t) \in [0, 2\pi] \times (0, T], \\ u(x, 0) &= \sin(x) - x, & x \in R \end{aligned}$$

with the periodic boundary condition, where the exact solution is

$$u(x, t) = e^{-t} \sin(x - t).$$

Table 6.1 Errors and orders for u . $\lambda = \theta, T = 1.0, k = 2, 3, 4$.

	N	$\ e_{un}\ $	Order	$\ e_u\ _c$	Order	$\ e_{u,r}\ $	Order	$\ e_{u,rx}\ $	Order
$k = 2$	20	5.20E-08	–	1.91E-07	–	4.53E-06	–	6.95E-05	–
$CFL = 0.01$	40	1.83E-09	4.83	6.23E-09	4.94	2.80E-07	4.01	8.74E-06	2.99
$\lambda = 0.8$	80	6.09E-11	4.91	1.99E-10	4.96	1.74E-08	4.01	1.10E-06	2.99
$\theta = 0.8$	160	1.96E-12	4.96	6.32E-12	4.98	1.08E-09	4.00	1.38E-07	2.99
$k = 3$	15	5.35E-10	–	6.62E-10	–	1.90E-07	–	1.27E-05	–
$CFL = 0.005$	30	3.82E-12	7.13	5.69E-12	6.86	5.53E-09	5.10	7.86E-07	4.02
$\lambda = 0.9$	45	2.07E-13	7.19	3.50E-13	6.88	7.12E-10	5.05	1.55E-07	4.00
$\theta = 0.9$	60	2.66E-14	7.13	4.80E-14	6.91	1.67E-10	5.01	4.89E-08	4.01
$k = 4$	10	1.60E-11	–	5.08E-11	–	8.34E-08	–	7.88E-06	–
$CFL = 0.001$	15	2.04E-13	10.76	1.23E-12	9.17	7.35E-09	5.98	1.06E-06	4.94
$\lambda = 1.2$	20	1.07E-14	10.23	7.91E-14	9.54	1.31E-09	6.01	2.54E-07	4.98
$\theta = 1.2$	25	6.74E-15	2.10	9.02E-15	9.73	3.41E-10	6.01	8.33E-08	4.99

Table 6.1 lists the results for u with $\lambda = \theta$, from which we observe $(2k + 1)$ th order superconvergence for numerical traces as well as cell averages, and that the convergence order of the error as well as its derivative are $k + 2$ and $k + 1$, respectively. **Table 6.2 shows errors and orders for q , demonstrating that our results hold true for the auxiliary variable when $\lambda = \theta$.** Moreover, the results with different weights

Table 6.2 Errors and orders for q . $\lambda = \theta, T = 1.0, k = 2, 3, 4$.

	N	$\ e_{qn}\ $	Order	$\ e_q\ _c$	Order	$\ e_{q,l}\ $	Order	$\ e_{q,lx}\ $	Order
$k = 2$	20	1.28E-07	–	2.50E-08	–	5.10E-06	–	8.50E-05	–
$CFL = 0.01$	40	4.15E-09	4.94	1.04E-09	4.59	3.19E-07	3.99	1.06E-05	3.00
$\lambda = 0.7$	80	1.33E-10	4.96	3.76E-11	4.79	2.00E-08	4.00	1.33E-06	2.99
$\theta = 0.7$	160	4.21E-12	4.98	1.26E-12	4.90	1.25E-09	4.00	1.67E-07	3.00
$k = 3$	15	1.55E-09	–	5.31E-10	–	4.14E-07	–	1.44E-05	–
$CFL = 0.005$	30	1.26E-11	6.94	3.81E-12	7.12	1.25E-08	5.05	8.92E-07	4.01
$\lambda = 0.9$	45	7.52E-13	6.96	2.07E-13	7.19	1.63E-09	5.02	1.76E-07	4.01
$\theta = 0.9$	60	1.02E-13	6.93	2.66E-14	7.12	3.83E-10	5.03	5.54E-08	4.01
$k = 4$	10	5.72E-11	–	1.58E-11	–	1.02E-07	–	8.12E-06	–
$CFL = 0.001$	15	1.60E-12	8.82	2.03E-13	10.74	8.79E-09	6.05	1.11E-06	4.91
$\lambda = 1.2$	20	1.32E-13	8.68	1.08E-14	10.21	1.55E-09	6.03	2.63E-07	5.00
$\theta = 1.2$	25	2.00E-14	8.45	6.80E-15	2.06	4.04E-10	6.03	8.70E-08	4.95

for λ, θ are given in Table 6.3, and similar conclusions can be observed for u , indicating that choosing different parameters for convection term and diffusion term does not affect the superconvergence results as far as the prime variable u is concerned.

Table 6.3 Errors and orders for u . $\lambda \neq \theta, T = 1.0, k = 2, 3, 4$.

	N	$\ e_{un}\ $	Order	$\ e_u\ _c$	Order	$\ e_{u,r}\ $	Order	$\ e_{u,rx}\ $	Order
$k = 2$	20	1.41E-07	–	3.09E-07	–	4.75E-06	–	6.71E-05	–
$CFL = 0.01$	40	4.60E-09	4.93	9.89E-09	4.97	2.91E-07	4.03	8.58E-06	2.97
$\lambda = 1.2$	80	1.47E-10	4.96	3.13E-10	4.98	1.80E-08	4.01	1.09E-06	2.97
$\theta = 0.8$	160	4.66E-12	4.98	9.88E-12	4.99	1.12E-09	4.01	1.37E-07	2.99
$k = 3$	15	1.85E-10	–	7.44E-10	–	2.59E-07	–	4.80E-06	–
$CFL = 0.002$	30	1.60E-12	6.85	5.46E-12	7.09	8.03E-09	5.01	3.01E-07	3.99
$\lambda = 0.9$	45	9.64E-14	6.93	3.13E-13	7.04	1.05E-09	5.01	5.96E-08	3.99
$\theta = 1.1$	60	1.28E-14	7.02	4.16E-14	7.02	2.50E-10	5.00	1.88E-08	4.00
$k = 4$	10	1.87E-10	–	1.69E-10	–	8.13E-08	–	7.88E-06	–
$CFL = 0.001$	15	4.93E-12	8.97	4.63E-12	8.88	7.20E-09	5.98	1.06E-06	4.95
$\lambda = 0.8$	20	3.54E-13	9.15	3.35E-13	9.12	1.28E-09	6.00	2.53E-07	4.98
$\theta = 1.2$	25	4.33E-14	9.41	4.09E-14	9.43	3.38E-10	5.97	8.31E-08	4.99

Example 6.2. We consider the following problem

$$\begin{aligned}
 u_t + u_x - u_{xx} &= 0, & (x, t) \in [0, 2\pi] \times (0, T], \\
 u(x, 0) &= \sin(x) - x, & x \in R
 \end{aligned}$$

with the mixed boundary conditions

$$u(0, t) = e^{-t} \sin(t) - t, \quad u_x(2\pi, t) = e^{-t} \cos(t) + 1;$$

Table 6.4 Errors and rates for mixed boundary condition (5.1).

N	$\lambda = \theta = 0.8$				$\lambda = \theta = 1.2$				
	$\ e_{un}\ $	Order	$\ e_u\ _c$	Order	$\ e_{un}\ $	Order	$\ e_u\ _c$	Order	
P^1	40	2.30E-05	–	3.48E-05	–	8.25E-06	–	1.35E-05	–
	80	2.72E-06	3.08	4.30E-06	3.02	1.06E-06	2.95	1.75E-06	2.94
	160	3.31E-07	3.04	5.35E-07	3.01	1.35E-07	2.97	2.24E-07	2.97
	320	4.07E-08	3.02	6.67E-08	3.00	1.71E-08	2.99	2.82E-08	2.98
P^2	20	7.36E-08	–	1.83E-07	–	4.49E-07	–	7.09E-07	–
	40	2.05E-09	5.16	5.64E-09	5.02	1.37E-08	5.04	2.24E-08	4.98
	80	6.10E-11	5.07	1.76E-10	5.00	4.16E-10	5.04	6.99E-10	5.00
	160	1.87E-12	5.04	5.10E-12	5.10	1.27E-11	5.03	2.15E-11	5.02
P^3	20	9.64E-11	–	1.56E-10	–	3.90E-11	–	8.65E-11	–
	30	5.53E-12	7.05	9.63E-12	6.88	2.26E-12	7.02	4.93E-12	7.07
	40	7.04E-13	7.16	1.32E-12	6.90	3.12E-13	6.88	6.45E-13	7.07
	50	1.65E-13	6.50	2.93E-13	6.77	9.82E-14	5.17	1.82E-13	5.66

the exact solution is

$$u(x, t) = e^{-t} \sin(x - t) + x - t.$$

The problem is solved by the LDG scheme (2.2) with $k = 1, 2, 3$, respectively, and the numerical fluxes are chosen as (5.2). We list various errors and corresponding convergence rates when $\lambda = \theta = 0.8, \lambda = \theta = 1.2$ in Table 6.4. The superconvergence results of order $2k + 1$ at numerical traces as well as cell averages demonstrate that the superconvergence also holds for mixed boundary conditions. In addition, to verify theoretical results for Dirichlet boundary conditions, we consider Example 6.2 with the following Dirichlet boundary conditions

$$u(0, t) = e^{-t} \sin(t) + t, \quad u(2\pi, t) = e^{-t} \cos(t) - 1.$$

The results are shown in Table 6.5, which confirms that the conclusion still holds for Dirichlet boundary conditions.

7 Concluding Remarks

In this paper, we obtain the superconvergence of the convection-diffusion equations based on the generalized alternating numerical fluxes. The main techniques are the construction of correction functions and analysis of the generalized Gauss–Radau projections and their modified versions. Different boundary conditions including periodic, mixed and Dirichlet boundary conditions are considered. The sharpness of the theoretical results is confirmed by numerical experiments. In further work, we will consider the degenerate diffusion problems and multidimensional equations.

Acknowledgements This work was supported by National Natural Science Foundation of China (Grants No. 11971132, U1637208, 71773024, 51605114, 11501149) and the National Key Research and Development Program of China (Grant No. 2017YFB1401801).

Table 6.5 Errors and rates for Dirichlet boundary condition (5.4).

N	$\lambda = \theta = 0.7$				$\lambda = \theta = 0.9$				
	$\ e_{un}\ $	Order	$\ e_u\ _c$	Order	$\ e_{un}\ $	Order	$\ e_u\ _c$	Order	
P^1	40	3.44E-05	-	5.27E-05	-	1.54E-05	-	2.51E-05	-
	80	4.15E-06	3.05	6.62E-06	2.99	1.95E-06	2.98	3.21E-06	2.96
	160	5.09E-07	3.03	8.27E-07	3.00	2.46E-07	2.99	4.07E-07	2.98
	320	6.28E-08	3.02	1.03E-07	3.00	3.09E-08	2.99	5.12E-08	2.99
P^2	20	2.03E-08	-	1.13E-07	-	7.07E-08	-	2.47E-07	-
	40	7.83E-10	4.69	3.69E-09	4.93	2.34E-09	4.92	7.89E-09	4.97
	80	3.13E-11	4.65	1.20E-10	4.94	7.56E-11	4.95	2.50E-10	4.98
	160	1.06E-12	4.88	3.81E-12	4.98	2.38E-12	4.99	7.83E-12	5.00
P^3	20	1.26E-10	-	2.05E-10	-	6.23E-11	-	1.25E-10	-
	30	7.77E-12	6.87	1.17E-11	7.06	3.78E-12	6.91	6.87E-12	7.15
	40	1.05E-12	6.94	1.65E-12	6.81	5.36E-13	6.79	1.02E-12	6.62
	50	2.57E-13	6.33	3.26E-13	7.26	1.98E-13	4.47	1.93E-13	7.48

References

- Ainsworth M, Monk P, Muniz W. Dispersive and dissipative properties of discontinuous Galerkin finite element methods for the second-order wave equation. *J Sci Comput*, 2006, 27: 5–40
- Bona J L, Chen H, Karakashian O, Xing Y. Conservative, discontinuous Galerkin–methods for the generalized Korteweg–de Vries equation. *Math Comput*, 2013, 82: 1401–1432
- Cao W, Li D, Yang Y, Zhang Z. Superconvergence of discontinuous Galerkin methods based on upwind-biased fluxes for 1D linear hyperbolic equations. *ESAIM: Math Model Numer Anal*, 2017, 51: 467–486
- Cao W, Liu H, Zhang Z. Superconvergence of the direct discontinuous Galerkin method for convection-diffusion equations. *Numer Methods for PDEs*, 2017, 33: 290–317
- Cao W, Shu C W, Yang Y, Zhang Z. Superconvergence of discontinuous Galerkin method for 2-D hyperbolic equations. *SIAM J Numer Anal*, 2015, 53: 1651–1671
- Cao W, Shu C W, Zhang Z. Superconvergence of discontinuous Galerkin methods for 1-D linear hyperbolic equations with degenerate variable coefficients. *ESAIM: Math Model Numer Anal*, 2017, 51: 2213–2235
- Cao W, Zhang Z. Superconvergence of local discontinuous Galerkin method for one-dimensional linear parabolic equations. *Math Comput*, 2016, 85: 63–84
- Cao W, Zhang Z, Zou Q. Superconvergence of discontinuous Galerkin method for linear hyperbolic equations. *SIAM J Numer Anal*, 2014, 52: 2555–2573
- Castillo P. An optimal error estimate for the local discontinuous Galerkin method. *Lectures Notes in Computational Science and Engineering*, 2000
- Castillo P, Cockburn B, Schötzau D, Schwab C. Optimal a priori error estimates for the hp -version of the local discontinuous Galerkin method for convection-diffusion problems. *Math Comput*, 2002, 71: 455–478
- Chen C, Hu S. The highest order superconvergence for bi - k degree rectangular elements at nodes a proof of $2k$ -conjecture. *Math Comput*, 2013, 82: 1337–1355
- Cheng Y, Chou C S, Li F, Xing Y. L^2 stable discontinuous Galerkin methods for one-dimensional two-way wave equations. *Math Comput*, 2017, 86: 121–155
- Cheng Y, Meng X, Zhang Q. Application of generalized Gauss–Radau projections for the local discontinuous Galerkin method for linear convection–diffusion equations. *Math Comput*, 2017, 86: 1233–1267
- Cheng Y, Shu C W. Superconvergence of discontinuous Galerkin and local discontinuous Galerkin schemes for linear hyperbolic and convection-diffusion equations in one space dimension. *SIAM J Numer Anal*, 2010, 47: 4044–4072
- Cockburn B, Luskin M, Shu C W, Suli E. Enhanced accuracy by post-processing for finite element methods for hyperbolic equations. *Math Comput*, 2003, 72: 577–606
- Cockburn B, Shu C W. The local discontinuous Galerkin method for time-dependent convection-diffusion systems. *SIAM J Numer Anal*, 1997, 35: 2440–2463

- 17 Docampo-S J, Ryan J K, Mirzargar M, Kirby R M. Multi-Dimensional Filtering: Reducing the Dimension Through Rotation. *SIAM J Sci Comput*, 2017, 39: A2179–A2200.
- 18 Freaun D, Ryan J. Superconvergence and the Numerical Flux: A study using the upwind-biased flux in discontinuous Galerkin methods. *Comm on Appl Math and Comput*, submitted
- 19 Fu P, Cheng Y, Li F, Xu Y. Discontinuous Galerkin methods with optimal L^2 accuracy for one dimensional linear PDEs with high order spatial derivatives. *J Sci Comput*, 2019, 78: 816–863
- 20 Hufford C, Xing Y. Superconvergence of the local discontinuous Galerkin method for the linearized Korteweg–de Vries equation. *J Comput and Appl Math*, 2014, 255: 441–455
- 21 Ji L, Xu Y, Ryan J. Accuracy-enhancement of discontinuous Galerkin solutions for convection-diffusion equations in multiple-dimensions. *Math Comput*, 2012, 81: 1929–1950
- 22 Liu H, Ploymaklam N. A local discontinuous Galerkin method for the Burgers–Poisson equation. *Numer Math*, 2015, 129: 321–351
- 23 Meng X, Shu C W, Wu B. Optimal error estimates for discontinuous Galerkin methods based on upwind-biased fluxes for linear hyperbolic equations. *Math Comput*, 2016, 85: 1225–1261
- 24 Meng X, Shu C W, Wu B. Superconvergence of the local discontinuous Galerkin method for linear fourth-order time-dependent problems in one space dimension. *IMA J Numer Anal*, 2012, 32: 1294–1328
- 25 Yang Y, Shu C W. Analysis of optimal superconvergence of local discontinuous Galerkin method for one-dimensional linear parabolic equations. *J Comput Math*, 2015, 33: 323–340